

Ionization of Atomic Hydrogen in Strong Infrared Laser Fields

Brant Abeln, Ethan Secor, Daniel Weflen, Klaus Bartschat, Timothy Urness, and Alexei N. Grum-Grzhimailo

Department of Physics and Astronomy, Drake University, Des Moines, IA 50311, USA

Department of Mathematics and Computer Science, Drake University, Des Moines, IA 50311, USA

Institute of Nuclear Physics, Moscow State University, Moscow 119991, Russia

Research Supported by the United States National Science Foundation


Abstract

We have used the matrix iteration method of Nurhuda and Faisal (Phys. Rev. A **60** (1999) 3125) to treat ionization of atomic hydrogen by a strong laser pulse. After testing our predictions against a variety of previous calculations, we present ejected-electron spectra as well as angular distributions for few-cycle infrared laser pulses with peak intensities of up to 10^{15} W/cm². The convergence of the results with the number of partial waves is a serious issue, which can be managed in a satisfactory way by using the velocity form of the electric dipole operator in connection with an efficient time-propagation scheme.

Introduction

- **1 attosecond** is one-millionth of one millionth of one millionth (10^{-18}) of a second.
- There are twice as many attoseconds in 1 second than seconds in the **age of the universe** (15 billion years)!
- The period for the $n = 1$ orbit in atomic hydrogen is about 150 attoseconds.
- Attosecond laser pulses provide a window to study the details of (valence) electron interactions in atoms and molecules.
- A major role for **theory** in attosecond science is to **suggest novel ways of investigating and controlling electronic processes** in matter on ultra-short time scales.
- Typical laser intensities in this field range from 10^{12} to 10^{15} W/cm².
- **10^{14} W/cm² is a million billion times stronger than the radiation that the Earth receives from the Sun directly above us on a clear day.**
- Such intensities can rip electrons away from atoms in very different ways from the **standard photoeffect**:
 - **Multi-photon ionization**
 - **Above-threshold ionization**
 - **Field (tunnel) ionization**

Single vs. Multi-Photon Ionization in Atomic Hydrogen



Numerical Method

- We start with the **Time-Dependent Schrödinger Equation**

$$\hat{H}\Psi = i\frac{\partial}{\partial t}\Psi \quad (1)$$

- In the **Length Form** of the electric dipole operator,

$$\hat{H} = -\frac{1}{2}\nabla^2 + \frac{l(l+1)}{2r^2} - \frac{1}{r} + r\cos(\vartheta)\mathbf{E}(t) \quad (2)$$

- In the **Velocity Form**, we have instead

$$\hat{H} = -\frac{1}{2}\nabla^2 + \frac{l(l+1)}{2r^2} - \frac{1}{r} - \frac{i\mathbf{A}(t)}{c} \cdot \nabla \quad (3)$$

- We propagate the initial wavefunction $\Psi(\mathbf{r}, t=0)$ in time using **Finite Differences**.

- In the **Crank-Nicholson Approximation**

$$\Psi(\mathbf{r}, t+\Delta t) \approx \frac{1-i\hat{H}\Delta t/2}{1+i\hat{H}\Delta t/2}\Psi(\mathbf{r}, t) \quad (4)$$

it is difficult to calculate the “denominator”, i.e., the inverse of a matrix at every time step.


Matrix Iteration Method

- We define $1+i\hat{H}\Delta t/2 \equiv \hat{O}_D + \hat{O}_{ND}$
- and write $(1+i\hat{H}\Delta t/2)^{-1} \approx (1-\hat{O}_D^{-1}\hat{O}_{ND} + \hat{O}_D^{-1}\hat{O}_{ND}\hat{O}_D^{-1}\hat{O}_{ND} + \dots)\hat{O}_D^{-1}$
- Taking 3–8 terms in the series expansion generally yields converged results.

Observables of Interest

- Observables are the **physical values** obtained from $\Psi(\mathbf{r}, t \rightarrow \infty)$ in a **measurement**.


Scheme of an Angular-Distribution Experiment




- **Photoelectron spectrum**: How many electrons come out, and with what energies?
- **Angular distribution of photoelectrons**: Where do the electrons go?

Results

Madroñero and Piraux (2009)



Our Calculation (2010, in press)



- The recent calculations by Madroñero and Piraux (Phys. Rev. A **80** (2009), 033409) exhibit two serious problems:
 - **unphysical oscillations**, most likely due to the insufficient size of their numerical grid;
 - **lack of convergence** for energies above ≈ 0.3 a.u., due to a limited number of basis functions.
- We pushed our calculations to even **higher intensities** (10^{15} W/cm²) **and energies** up to 4 a.u.
- **Measurements are currently in progress** at Griffith University in Brisbane (Australia).
- **We generated movies to illustrate the time dependence of the ionization process.**
- Below are snapshots for a 4-cycle pulse with a \sin^2 envelope function, a maximum intensity of 10^{15} W/cm², and a central wavelength of 152 nm.

